



On Binary Classification in Extreme Regions

Hamid Jalalzai, S. Cléménçon, Anne Sabourin

► **To cite this version:**

Hamid Jalalzai, S. Cléménçon, Anne Sabourin. On Binary Classification in Extreme Regions. NIPS, Sep 2018, Montréal, Canada. hal-01932813

HAL Id: hal-01932813

<https://hal-imt.archives-ouvertes.fr/hal-01932813>

Submitted on 23 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On Binary Classification in Extreme Regions

Hamid Jalalzai, Stephan Cléménçon and Anne Sabourin

LTCI Telecom ParisTech, Université Paris-Saclay
75013, Paris, France
first.last@telecom-paristech.fr

Abstract

In pattern recognition, a random label Y is to be predicted based upon observing a random vector X valued in \mathbb{R}^d with $d \geq 1$ by means of a classification rule with minimum probability of error. In a wide variety of applications, ranging from finance/insurance to environmental sciences through teletraffic data analysis for instance, extreme (*i.e.* very large) observations X are of crucial importance, while contributing in a negligible manner to the (empirical) error however, simply because of their rarity. As a consequence, empirical risk minimizers generally perform very poorly in extreme regions. It is the purpose of this paper to develop a general framework for classification in the extremes. Precisely, under non-parametric heavy-tail assumptions for the class distributions, we prove that a natural and asymptotic notion of risk, accounting for predictive performance in extreme regions of the input space, can be defined and show that minimizers of an empirical version of a non-asymptotic approximant of this dedicated risk, based on a fraction of the largest observations, lead to classification rules with good generalization capacity, by means of maximal deviation inequalities in low probability regions. Beyond theoretical results, numerical experiments are presented in order to illustrate the relevance of the approach developed.

1 Introduction

Because it covers a wide range of practical applications and its probabilistic theory can be straightforwardly extended to some extent to various other prediction problems, binary classification can be considered as the flagship problem in statistical learning. In the standard setup, (X, Y) is a random pair defined on a certain probability space with (unknown) joint probability distribution P , where the (output) r.v. Y is a binary label, taking its values in $\{-1, +1\}$ say, and X models some information, valued in \mathbb{R}^d and hopefully useful to predict Y . In this context, the goal pursued is generally to build, from a training sample $\mathcal{D}_n = \{(X_1, Y_1), \dots, (X_n, Y_n)\}$ composed of $n \geq 1$ i.i.d. realizations of (X, Y) , a classifier $g : \mathbb{R}^d \rightarrow \{-1, +1\}$ minimizing the probability of error $L_P(g) = \mathbb{P}\{Y \neq g(X)\}$. The Empirical Risk Minimization paradigm (ERM in abbreviated form, see *e.g.* [6]) suggests considering solutions g_n of the minimization problem $\min_{g \in \mathcal{G}} \widehat{L}_n(g)$, where $\widehat{L}_n(g)$ is a statistical estimate of the risk $L(g)$. In general the empirical version $\widehat{L}_n(g) = (1/n) \sum_{i=1}^n \mathbb{1}\{Y_i \neq g(X_i)\}$ is considered, denoting by $\mathbb{1}\{\mathcal{E}\}$ the indicator function of any event \mathcal{E} . This amounts to replacing P in L_P with the empirical distribution of the (X_i, Y_i) 's. The class \mathcal{G} of predictive rules is supposed to be rich enough to contain a reasonable approximant of the minimizer of L_P , *i.e.* the Bayes classifier $g^*(x) = 2 \mathbb{1}\{\eta(x) \geq 1/2\} - 1$, where $\eta(X) = \mathbb{P}\{Y = 1 \mid X\}$ denotes the posterior probability.

Because extreme observations X , *i.e.* observations whose norm $\|X\|$ exceeds some large threshold $t > 0$, are rare and thus underrepresented in the training dataset \mathcal{D}_n classification errors in these

regions of the input space may have a negligible impact on the global prediction error of \hat{g}_n . Notice incidentally that the threshold t may depend on n , since ‘large’ should be naturally understood as large w.r.t the vast majority of data previously observed. Using the total probability formula, one may indeed write

$$L_P(g) = \mathbb{P}\{\|X\| > t\}\mathbb{P}\{Y \neq g(X) \mid \|X\| > t\} + \mathbb{P}\{\|X\| \leq t\}\mathbb{P}\{Y \neq g(X) \mid \|X\| \leq t\}. \quad (1)$$

Hence, due to the extremely small order of magnitude of $\mathbb{P}\{\|X\| > t\}$ and of its empirical counterpart, there is no guarantee that the standard ERM strategy produces an optimal classifier on the extreme region $\{x : \|x\| > t\}$. In other words the quantity $\mathbb{P}\{Y \neq \hat{g}_n(X) \mid \|X\| > t\}$ may not be nearly optimal, whereas in certain practical applications (*e.g.* finance, insurance, environmental sciences, aeronautics safety), accurate prediction in extreme regions is crucial.

The purpose of the subsequent analysis is to investigate the problem of building a classifier such that the first term of the decomposition (1) is asymptotically minimum as $t \rightarrow +\infty$. We thus consider the conditional probability of error, which quantity is next referred to as the *classification risk above level t* , given by

$$L_t(g) := L_{P_t}(g) = \mathbb{P}\{Y \neq g(X) \mid \|X\| > t\}, \quad (2)$$

denoting by P_t the conditional distribution of (X, Y) given $\|X\| > t$. In this paper, we address the issue of learning a classifier g_n whose risk $L_t(g_n)$ is asymptotically minimum as $t \rightarrow \infty$ with high probability. In order to develop a framework showing that a variant of the ERM principle tailored to this statistical problem leads to predictive rules with good generalization capacities, (non-parametric) distributional assumptions related to the tail behavior of the class distributions F_+ and F_- , the conditional distributions of the input r.v. X given $Y = +/ - 1$, are required. Precisely, we assume that they are both multivariate regularly varying, which correspond to a large non-parametric class of (heavy-tailed) distributions, widely used in applications where the impact of extreme observations should be enhanced, or at least not neglected. Hence, under appropriate non-parametric assumptions on F_+ and F_- , as well as on the tail behavior of $\eta(x)$, we prove that $\min_g L_t(g)$ converges to a quantity denoted by L_∞^* and referred to as the *asymptotic risk in the extremes*, as $t \rightarrow \infty$. It is also shown that this limit can be interpreted as the minimum classification error related to a (non observable) random pair (X_∞, Y_∞) , whose distribution P_∞ corresponds to the limit of the conditional distribution of (X, Y) given $\|X\| > t$, for an appropriate normalization of X , as $t \rightarrow \infty$. With respect to the goal set above we next investigate the performance of minimizer $\hat{g}_{n,\tau}$ of an empirical version of the risk $L_{P_{t_\tau}}$, where t_τ is the $(1 - \tau)$ quantile of the r.v. $\|X\|$ and $\tau \ll 1$. The computation of $\hat{g}_{n,\tau}$ involves the $\lfloor n\tau \rfloor$ input observations with largest norm, and the minimization is performed over a collection of classifiers of finite VC dimension. Based on a variant of the VC inequality tailored to low probability regions, rate bounds for the deviation $L_t(\hat{g}_{n,\tau}) - L_\infty^*$ are established, of order $O_{\mathbb{P}}(1/\sqrt{n\tau})$ namely. These theoretical results are also illustrated by preliminary experiments based on synthetic data.

The rest of the paper is organized as follows. Multivariate extreme value theory (MEVT) notions involved in the framework we develop are described in section 2, together with the probabilistic setup we consider for classification in the extremes. A notion of risk tailored to this statistical learning task is also introduced therein. Section 3 investigates how to extend the ERM principle in this situation. In particular, probability bounds proving the generalization ability of minimizers of a non-asymptotic approximant of the risk previously introduced are established. Illustrative numerical results are displayed in section 4, while several concluding remarks are collected in section 5. Some technical details and proofs are deferred to the Supplementary Material.

2 Probabilistic Framework - Preliminary Results

We start off with recalling concepts pertaining to MEVT and next develop a general framework in order to formulate the problem of binary classification in the extremes in a rigorous manner. For completeness, additional details about regular variation and vague convergence are given in the supplementary material (Appendix A).

2.1 Regularly Varying Random Vector

By definition, heavy-tail phenomena are those which are ruled by very large values, occurring with a far from negligible probability and with significant impact on the system under study. When the

phenomenon of interest is described by the distribution of a univariate random variable, the theory of regularly varying functions provides the appropriate mathematical framework for the study of heavy-tailed distributions. One may refer to [13] for an excellent account of the theory of regularly varying functions and its application to the study of heavy-tailed distributions. For examples of works where such assumptions are considered in the context of statistical learning, see *e.g.* [8, 3, 14, 12, 1] or [10]. Let $\alpha > 0$, a random variable X is said to be regularly varying with tail index α if

$$\mathbb{P}\{X > tx \mid X > t\} \xrightarrow[t \rightarrow \infty]{} x^{-\alpha}, \quad x > 1.$$

This is the case if and only if there exists a function $b : \mathbb{R}_+ \rightarrow \mathbb{R}_+^*$ with $b(t) \rightarrow \infty$ such that for all $x > 0$, the quantity $t\mathbb{P}\{X/b(t) > x\}$ converges to some limit $h(x)$ as $t \rightarrow \infty$. Then b may be chosen as $b(t) = t^{1/\alpha}$ and $h(x) = cx^{-\alpha}$ for some $c > 0$. Based on this characterization, the heavy-tail model can be extended to the multivariate setup. Consider a d -dimensional random vector $X = (X^{(1)}, \dots, X^{(d)})$ taking its values in \mathbb{R}_+^d . Assume that all the $X^{(j)}$ are regularly varying with index $\alpha > 0$. Then the random *vector* X is said to be regularly varying with tail index α if there exists a non null positive Radon measure μ on the punctured space $E = [0, \infty]^d \setminus \{0\}$ and a function $b(t) \rightarrow \infty$ such that for all Borel set $A \subset E$ such that $0 \notin \partial A$ and $\mu(\partial A) = 0$,

$$t\mathbb{P}\{X/b(t) \in A\} \xrightarrow[t \rightarrow \infty]{} \mu(A).$$

In such a case, the so-called *exponent measure* μ fulfills the homogeneity property $\mu(tC) = t^{-\alpha}\mu(C)$ for all $t > 0$ and any Borel set $C \subset E$. This suggests a decomposition of μ into a radial component and an angular component Φ . For all $x = (x_1, \dots, x_d) \in \mathbb{R}_+^d$, set

$$\begin{cases} R(x) = \|x\|, \\ \Theta(x) = \left(\frac{x_1}{R(x)}, \dots, \frac{x_d}{R(x)} \right) \in S, \end{cases}$$

where S is the positive orthant of the unit sphere in \mathbb{R}^d for the chosen norm $\|\cdot\|$. The choice of the norm is unimportant as all norms are equivalent in \mathbb{R}^d . Define an *angular measure* Φ on S as

$$\Phi(B) = \mu\{r\theta : \theta \in B, r \geq 1\}, \quad B \subset S, \text{ measurable.}$$

The angular measure Φ is finite, and the conditional distribution of $(R(X)/t, \Theta(X))$ given that $R(X) > t$ converges as $t \rightarrow \infty$ to a limit which admits the following product decomposition: for $r \geq 1$ and $B \subset S$ such that $\Phi(\partial B) = 0$,

$$\begin{aligned} \lim_{t \rightarrow \infty} \mathbb{P}\{R(X)/t > r, \Theta(X) \in B \mid R(X) > t\} &= c \mu\{x : R(x) > r, \Theta(x) \in B\} \\ &= c r^{-\alpha} \Phi(B), \end{aligned}$$

where $c = \mu\{x : R(x) > 1\}^{-1} = \Phi(S)^{-1}$ is a normalizing constant. Thus $c\Phi$ may be viewed as the limiting distribution of $\Theta(X)$ given that $R(X)$ is large.

Remark 1 *It is assumed above that all marginal distributions are tail equivalent to the Pareto distribution with index α . In practice, the tails of the marginals may be different and it may be convenient to work with marginally standardized variables, that is, to separate the margins $F_j(x_j) = \mathbb{P}\{X^{(j)} \leq x_j\}$ from the dependence structure in the description of the joint distribution of X . Consider the standardized variables $V^{(j)} = 1/(1 - F_j(X^{(j)})) \in [1, \infty]$ and $V = (V^{(1)}, \dots, V^{(d)})$. Replacing X by V permits to take $\alpha = 1$ and $b(t) = t$.*

2.2 Classification in the Extremes - Assumptions, Criterion and Optimal Elements

We place ourselves in the binary classification framework recalled in the introduction. For simplicity, we suppose that X takes its values in the positive orthant \mathbb{R}_+^d . The general aim is to build from training data in the extreme region (*i.e.* data points (X_i, Y_i) such that $\|X_i\| > t_n$ for a large threshold value $t_n > 0$) a classifier $g_n(x)$ with risk $L_{t_n}(g_n)$ defined in (2) being asymptotically minimum as $t_n \rightarrow \infty$. In this purpose, we introduce general assumptions guaranteeing that the minimum risk $L_t(g^*)$ above level t has a limit as $t \rightarrow \infty$. Throughout the article, we assume that the class distributions F_+ and F_- are heavy-tailed with same index $\alpha = 1$.

Assumption 1 For all $\sigma \in \{-, +\}$, the conditional distribution of X given $Y = \sigma 1$ is regularly varying with index 1 and angular measure $\Phi_\sigma(d\theta)$ (respectively, exponent measure $\mu_\sigma(dx)$): for $A \subset [0, \infty]^d \setminus \{0\}$ a measurable set such that $0 \notin \partial A$ and $\mu(\partial A) \neq 0$,

$$t\mathbb{P}\{t^{-1}X \in A \mid Y = \sigma 1\} \xrightarrow{t \rightarrow \infty} \mu_\sigma(A), \quad \sigma \in \{-, +\},$$

and for $B \subset S$ a measurable set,

$$\Phi_\sigma(B) = \mu_\sigma\{x \in \mathbb{R}_+^d : R(x) > 1, \Theta(x) \in B\}, \quad \sigma \in \{-, +\}.$$

Under the hypothesis above, X 's marginal distribution, given by $F = pF_+ + (1-p)F_-$, where $p = \mathbb{P}\{Y = +1\} > 0$, is heavy-tailed as well with index 1. Indeed, we have:

$$t\mathbb{P}\{t^{-1}X \in A\} \xrightarrow{t \rightarrow \infty} \mu(A) := p\mu_+(A) + (1-p)\mu_-(A).$$

And similarly

$$\Phi(B) := p\Phi_+(B) + (1-p)\Phi_-(B).$$

Observe also that the limiting class balance can be expressed using the latter asymptotic measures. Indeed, let $\Omega = \{x \in \mathbb{R}_+^d : \|x\| \leq 1\}$ denote the positive orthant of the unit ball and let Ω^c denote its complementary set in \mathbb{R}_+^d . We have:

$$p_t = \mathbb{P}\{Y = +1 \mid \|X\| > t\} = \frac{t\mathbb{P}\{\|X\| > t \mid Y = 1\}p}{t\mathbb{P}\{\|X\| > t\}} \xrightarrow{t \rightarrow \infty} p \frac{\mu_+(\Omega^c)}{\mu(\Omega^c)} = p \frac{\Phi_+(S)}{\Phi(S)} \stackrel{\text{def}}{=} p_\infty. \quad (3)$$

Remark 2 (ON ASSUMPTION 1) We point out that only the situation where the supposedly heavy-tailed class distributions F_+ and F_- have the same tail index is of interest. Suppose for instance that the tail index α_+ of F_+ is strictly larger than that of F_- , α_- , that is F_- has heavier tail than F_+ . In such a case F is still regularly varying with index $\min\{\alpha_+, \alpha_-\}$ and $p_t \rightarrow 0$. In this case, one may straightforwardly see that the classifier predicting always -1 on $\{x \in \mathbb{R}_+^d : \|x\| > t\}$ is optimal as t increases to infinity.

Remark 3 (ON ASSUMPTION 1 (BIS)) As noticed in Remark 1, assuming that $\alpha = 1$ is not restrictive when the marginal distributions are known. In practice however, they must be estimated. Due to space limitations, in the present analysis, we shall neglect the estimation error arising from their estimation. Relaxing this assumption, as made in e.g. [9], will be the subject of future work.

Asymptotic criterion for classification in the extremes. The goal pursued is to construct a classifier g_n , based on the training examples \mathcal{D}_n , minimizing the *asymptotic risk in the extremes* given by

$$L_\infty(g) = \limsup_{t \rightarrow \infty} L_t(g). \quad (4)$$

We also set $L_\infty^* = \inf_{g \text{ measurable}} L_\infty(g)$. It is immediate that any classifier which coincides with the Bayes classifier g^* on the region $t\Omega^c = \{x \in \mathbb{R}_+^d : \|x\| > t\}$ is optimal w.r.t. the distribution P_t . In particular g^* minimizes L_t and the associated risk is

$$L_t^* := L_t(g^*) = \mathbb{E}[\min\{\eta(X), 1 - \eta(X)\} \mid \|X\| > t], \quad t > 0. \quad (5)$$

Thus, for all classifier g , $L_t(g) \geq L_t(g^*)$, and taking the limit superior shows that g^* minimizes L_∞ , that is $L_\infty^* = L_\infty(g^*)$.

Optimality. The objective formulated above can be connected with a standard binary classification problem, related to a random pair (X_∞, Y_∞) taking its values in the limit space $\Omega^c \times \{-1, +1\}$, see Theorem 1 below. Let $\mathbb{P}\{Y_\infty = +1\} = p_\infty$ as in (3) and define the distribution of X_∞ given that $Y_\infty = \sigma 1$, $\sigma \in \{-, +\}$ as $\mu_\sigma(\Omega^c)^{-1}\mu_\sigma(\cdot)$. Then for $A \subset \Omega^c$, using (3),

$$\begin{aligned} \mathbb{P}\{X_\infty \in A, Y_\infty = +1\} &= \frac{p_\infty \mu_+(A)}{\mu_+(\Omega^c)} = \frac{p \mu_+(A)}{\mu(\Omega^c)} = \frac{p \lim_t t\mathbb{P}\{X \in tA \mid Y = +1\}}{\lim_t t\mathbb{P}\{X \in t\Omega^c\}} \\ &= \lim_{t \rightarrow \infty} \mathbb{P}\{X \in tA, Y = +1 \mid \|X\| > t\}. \end{aligned}$$

We denote by P_∞ the joint distribution of (X_∞, Y_∞) thus defined. As shall be seen below, under appropriate and natural assumptions, classifiers with minimum asymptotic risk in the extremes are in 1-to-1 correspondence with solutions of the binary classification problem related to (X_∞, Y_∞) . Let ρ be a common dominating measure for Φ_-, Φ_+ on S (ρ does not need to be the Lebesgue measure, take e.g. $\rho = \Phi_+ + \Phi_-$). Then denote by φ_+, φ_- respectively the densities of Φ_+, Φ_- w.r.t. ρ . By homogeneity of μ_+, μ_- , the conditional distribution of Y_∞ given $X_\infty = x$ is

$$\begin{aligned} \eta_\infty(x) &\stackrel{\text{def}}{=} \mathbb{P}\{Y_\infty = 1 \mid X_\infty = x\} = \frac{p_\infty \varphi_+(\Theta(x)) / \Phi_+(S)}{p_\infty \varphi_+(\Theta(x)) / \Phi_+(S) + (1 - p_\infty) \varphi_-(\Theta(x)) / \Phi_-(S)} \\ &= \frac{p \varphi_+(\Theta(x))}{p \varphi_+(\Theta(x)) + (1 - p) \varphi_-(\Theta(x))}. \end{aligned}$$

Notice that η_∞ is independent of the chosen reference measure ρ and that η_∞ is constant along rays, that is $\eta_\infty(tx) = \eta_\infty(x)$ for (t, x) such that $\min(\|tx\|, \|x\|) \geq 1$. The optimal classifier for the random pair (X_∞, Y_∞) with respect to the classical risk L_{P_∞} is clearly

$$g_\infty^*(x) = 2\mathbb{1}\{\eta_\infty(x) \geq 1/2\} - 1.$$

Again g_∞^* is constant along rays on Ω^c and is thus a function of $\Theta(x)$ only. We abusively denote $\eta_\infty(x) = \eta_\infty(\Theta(x))$. The minimum classification error is

$$L_{P_\infty}^* = L_{P_\infty}(g_\infty^*) = \mathbb{E}[\min\{\eta_\infty(\Theta_\infty), 1 - \eta_\infty(\Theta_\infty)\}], \quad (6)$$

where $\Theta_\infty = \Theta(X_\infty)$. More generally, observe that any class \mathcal{G}_S of classifiers $g : \theta \in S \mapsto g(\theta) \in \{-1, +1\}$ defines a class of classifiers on \mathbb{R}_+^d , $x \in \mathbb{R}_+^d \mapsto g(\Theta(x))$, that shall still be denoted by \mathcal{G}_S for simplicity. The next result claims that, under the regularity hypothesis stated below, the classifier g_∞^* is optimal for the asymptotic risk in the extremes, that is $L_\infty(g_\infty^*) = \inf_g L_\infty(g)$. We shall also prove that $L_\infty(g_\infty^*) = L_{P_\infty}^*$.

Assumption 2 (UNIFORM CONVERGENCE ON THE SPHERE OF $\eta(tx)$) *The limiting regression function η_∞ is continuous on S and*

$$\sup_{\theta \in S} |\eta(\Theta(t\theta)) - \eta_\infty(\theta)| \xrightarrow{t \rightarrow \infty} 0$$

Remark 4 (ON ASSUMPTION 2) *By invariance of η_∞ along rays, Assumption 2 is equivalent to*

$$\sup_{\{x \in \mathbb{R}_+^d : \|x\| \geq t\}} |\eta(x) - \eta_\infty(x)| \xrightarrow{t \rightarrow \infty} 0.$$

Assumption 2 is satisfied whenever the probability densities f_+, f_- of F_+, F_- are continuous, regularly varying with limit functions q_+, q_- , and when the convergence is uniform, that is if

$$\lim_{t \rightarrow \infty} \sup_{x \in S} |t^{d+1} f_\sigma(tx) - q_\sigma(x)| = 0, \quad \sigma \in \{+, -\}. \quad (7)$$

In such a case q_+, q_- are respectively the densities of μ_+, μ_- with respect to the Lebesgue measure and are continuous, which implies the continuity of φ_+, φ_- . The latter uniform convergence assumption is introduced in [5] and is used e.g. in [2] in the context of minimum level sets estimation.

Theorem 1 (OPTIMAL CLASSIFIERS IN THE EXTREMES) *Under Assumptions 1 and 2,*

$$L_t^* \xrightarrow{t \rightarrow \infty} L_{P_\infty}^*. \quad (8)$$

Hence, we have: $L_\infty^ = L_{P_\infty}^*$. In addition, the classifier g_∞^* minimizes the asymptotic risk in the extremes:*

$$\inf_{g \text{ measurable}} L_\infty(g) = L_\infty(g_\infty^*) = \mathbb{E}\{\min(\eta_\infty(\Theta_\infty), 1 - \eta_\infty(\Theta_\infty))\}.$$

Refer to the Supplementary Material for the technical proof. Theorem 1 gives us the form of the optimal classifier in the extremes $g_\infty^*(x) = g_\infty^*(\Theta(x))$, which depends only on the angular component $\Theta(x)$, not the norm $R(x)$. This naturally leads to applying the ERM principle to a collection of classifiers of the form $g(x) = g(\Theta(x))$ on the domain $\{x \in \mathbb{R}_+^d : \|x\| > t\}$ for $t > 0$ large enough. The next section provides statistical guarantees for this approach.

3 Empirical Risk Minimization in the Extremes

Consider a class \mathcal{G}_S of classifiers $g : \theta \in S \mapsto g(\theta) \in \{-1, +1\}$ on the sphere S . It also defines a collection of classifiers on \mathbb{R}_+^d , namely $\{g(\Theta(x)) : g \in \mathcal{G}_S\}$, which we denote by \mathcal{G}_S for simplicity. Sorting the training observations by decreasing order of magnitude, we introduce the order statistics $\|X_{(1)}\| > \dots > \|X_{(n)}\|$ and we denote by $Y_{(i)}$ the corresponding sorted labels. Fix a small fraction $\tau > 0$ of extreme observations, and let t_τ be the quantile at level $(1 - \tau)$ of the r.v. $\|X\|$: $\mathbb{P}\{\|X\| > t_\tau\} = \tau$. Set $k = \lfloor n\tau \rfloor$ and consider the empirical risk

$$\widehat{L}_k(g) = \frac{1}{k} \sum_{i=1}^k \mathbf{1}\{Y_{(i)} \neq g(\Theta(X_{(i)}))\} = L_{\widehat{P}_k}(g), \quad (9)$$

where \widehat{P}_k denotes the empirical distribution of the truncated training sample $\{(X_i, Y_i) : \|X_i\| \geq \|X_k\|, i \in \{1, \dots, n\}\}$, the statistical version of the conditional distribution P_{t_τ} . We now investigate the performance in terms of asymptotic risk in the extremes L_∞ of the solutions of the minimization problem

$$\min_{g \in \mathcal{G}_S} \widehat{L}_k(g). \quad (10)$$

The practical issue of designing efficient algorithms for solving (10) is beyond the scope of this paper. Focus is here on the study of the learning principle that consists in assigning to any very large input value x the likeliest label based on the direction $\Theta(x)$ it defines only (the construction is summarized in Algorithm 1 below). The following result provides an upper bound for the excess of classification error in the domain $t_\tau\Omega^c$ of solutions of (10). Its proof, which relies on a maximal deviation inequality tailored to low probability regions, is given in the Supplementary Material.

Theorem 2 *Suppose that the class \mathcal{G}_S is of finite VC dimension $V_{\mathcal{G}_S} < +\infty$. Let \widehat{g}_k be any solution of (10). Recall $k = \lfloor n\tau \rfloor$. Then, for $\delta \in (0, 1)$, $\forall n \geq 1$, we have with probability larger than $1 - \delta$:*

$$\begin{aligned} L_{t_\tau}(\widehat{g}_k) - L_{t_\tau}^* &\leq \frac{1}{\sqrt{k}} \left(\sqrt{2(1-\tau) \log(2/\delta)} + C\sqrt{V_{\mathcal{G}_S} \log(1/\delta)} \right) \\ &\quad + \frac{1}{k} \left(5 + 2 \log(1/\delta) + \sqrt{\log(1/\delta)} (C\sqrt{V_{\mathcal{G}_S}} + \sqrt{2}) \right) + \left\{ \inf_{g \in \mathcal{G}_S} L_{t_\tau}(g) - L_{t_\tau}^* \right\}, \end{aligned}$$

where C is a constant independent from n , τ and δ .

Remark 5 (ON MODEL SELECTION) *Selecting an appropriate model class \mathcal{G}_S is a crucial issue in machine-learning. Following in the footsteps of structured risk minimization, one may use a VC bound for $\mathbb{E}[\sup_{g \in \mathcal{G}_S} |\widehat{L}_k(g) - \mathbb{E}[\widehat{L}_k(g)]|]$ as a complexity regularization term to penalize in an additive fashion the empirical risk (9). Considering a collection of such models, oracle inequalities guaranteeing the quasi-optimality of the rule minimizing the penalized empirical risk can be then classically established by means of a slight modification of the argument of Theorem 2's proof, see e.g. Chapter 18 in [6].*

The upper bound stated above shows that the learning rate is of order $O_{\mathbb{P}}(1/\sqrt{k})$, where k is the actual size of the training data set used to perform approximate empirical risk minimization in the extremes. As revealed by the corollary below, this strategy permits to build a consistent sequence of classifiers for the L_∞ -risk, when the fraction $\tau = \tau_n$ decays at an appropriate rate (provided that the model bias can be neglected of course).

Corollary 1 *Suppose that the assumptions of Theorems 1-2 are fulfilled. In addition, assume that the model bias asymptotically vanishes as $\tau \rightarrow 0$, i.e.*

$$\inf_{g \in \mathcal{G}_S} L_{t_\tau}(g) - L_{t_\tau}^* \longrightarrow 0 \quad \text{as } \tau \rightarrow 0.$$

Then, as soon as $k \rightarrow +\infty$ as $n \rightarrow \infty$, the sequence of classifiers (\widehat{g}_k) is consistent in the extremes, meaning that we have the convergence in probability:

$$L_\infty(\widehat{g}_k) \rightarrow L_\infty^* \text{ as } n \rightarrow \infty.$$

Algorithm 1 (ERM in the extremes)

Input Training dataset $\mathcal{D}_n = \{(X_1, Y_1), \dots, (X_n, Y_n)\}$, collection \mathcal{G}_S of classifiers on the sphere, size $k \leq n$ of the training set composed of extreme observations

- 1 **Standardization.** Standardize the input vector by applying the rank-transformation: $\forall i \in \{1, \dots, n\}$, $\hat{V}_i = \hat{T}(X_i)$, where

$$\hat{T}(x) = \left(1 / \left(1 - \hat{F}_j(x_j)\right)\right)_{j=1, \dots, d},$$

for all $x = (x_1, \dots, x_d) \in \mathbb{R}^d$.

- 2 **Truncation.** Sort the training input observations by decreasing order of magnitude

$$\|\hat{V}_{(1)}\| \geq \dots \geq \|\hat{V}_{(n)}\|,$$

and consider the set of extreme training points

$$\left\{(\hat{V}_{(1)}, Y_{(1)}), \dots, (\hat{V}_{(k)}, Y_{(k)})\right\}.$$

- 3 **Optimization.** Compute a solution $\hat{g}_k(\theta)$ of the minimization problem

$$\min_{g \in \mathcal{G}_S} \frac{1}{k} \sum_{i=1}^k \mathbf{1} \left\{ Y_{(i)} \neq g \left(\Theta(\hat{V}_{(i)}) \right) \right\}$$

Output The classifier $\hat{g}_k \left(\Theta \left(\hat{T}(x) \right) \right)$, applicable on the region $\{x : \|\hat{T}(x)\| > \|\hat{V}_{(k)}\|\}$.

Remark 6 (Choice of k) Determining the best value of k is a typical challenge of Extreme Value analysis. This is typically a bias/variance trade-off, too large values introduce a bias by taking into account observations which are not large enough, so that their distribution deviates significantly from the limit distribution of extremes. On the other hand, too small values obviously increase the variance of the classifier. See e.g. [8] or [9] and the reference therein for a discussion. In practice a possible default choice is $k = \sqrt{n}$, otherwise cross-validation can be performed.

4 Illustrative Numerical Experiments

The purpose of our experiments is to provide insights into the performance of the classifier \hat{g}_k on extreme regions constructed via Algorithm 1. The training set is ordered as in Step 1 of Algorithm 1. For a chosen k , let $t = \|\hat{T}(X_{(k)}^{\text{train}})\|$, the L_1 norm is used throughout our experiments. The extreme test set \mathcal{T} is the subset of test points such that $\|\hat{T}(X_i^{\text{test}})\| > t$. To approximate of the asymptotic risk in the extremes $L_\infty(\hat{g}_k)$ and illustrate the generalization ability of the proposed classifier in the extreme region, we consider decreasing subsets of \mathcal{T} . Namely denoting $n_{\text{test}} = |\mathcal{T}|$, we keep only the $\lfloor \kappa n_{\text{test}} \rfloor$ largest instances of \mathcal{T} in terms on $\|\hat{T}(X_i^{\text{test}})\|$, for decreasing values of $\kappa \in (0, 1]$. This experimental framework is summarized in Figure 1, where $\lambda t = \|\hat{T}(X_{(\lfloor \kappa n_{\text{test}} \rfloor)}^{\text{test}})\| \geq t$.

We consider two different classification algorithms for Step 3 in Algorithm 1, namely random forest (RF) and k-nearest neighbors (k-NN), which correspond to two different classes \mathcal{G}_S of classifiers. For each class \mathcal{G}_S , the performance of \hat{g}_k (which considers only the direction $\Theta(\hat{T}(x))$ of both training and testing data, in other words classifies the projected datasets onto the unit sphere (see Figure 2) is compared with that of the classical version of the algorithm (RF or k-NN) taking as input the same training data but without the standardization and truncation steps neither the projection onto the unit sphere. Figures 4 and 5 summarize the results obtained using RF respectively with a multivariate simulated dataset and with a real world dataset. The simulated dataset is generated from a logistic distribution as described in [15]. The positive and negative instances are generated using




Figure 1: Train set (dotted area) and test set (colored area).




Figure 2: Colored cones correspond to a given label from the classifier on the simplex.

two different dependency parameters. An example of dataset thus obtained is displayed in Figure 3. We report the results obtained with $5 \cdot 10^3$ points for each label for the train set and $5 \cdot 10^4$ points for each label for the test set. $k = 100$ and $\kappa \in [1, 0.3]$. the number of trees for both random forests (in the regular setting and in the setting of Algorithm 1) is set to 200. The number of neighbors for both k-NN's is set to 5.




Figure 3: Toy dataset generated from a multivariate logistic distribution projected on \mathbb{R}^2 .

The real dataset known as Ecoli dataset, introduced in [11], deals with protein localization and contains 336 instances and 8 features. The Supplementary Material gathers additional details concerning the datasets and the tuning of RF and k-NN in our experiments, as well as additional results obtained with the above described datasets and with a simulated dataset from a different distribution.




Figure 4: Logistic data - test loss of RF on the simplex and regular RF depending on the multiplicative factor κ .




Figure 5: Real data - test loss of RF on the simplex and regular RF depending on the multiplicative factor κ .

Figure 4 shows the evolutions of the Hamming losses with decreasing values of $\kappa \in [0.3, 1]$. The boxplots display the losses obtained with 10 independently simulated datasets. For the experiment on the Ecoli dataset (Figure 5), one third of the dataset is used as a test set and the rest corresponds to the train set. $k = 100$ and $\kappa \in [0.3, 1]$ (considering smaller values of κ was prevented by data scarcity). The boxplots display the results for different (random) partitions of the data into a train and a test set. In both examples, the loss of the regular classifier is worse (and even increases) when κ decreases whereas the classifier resulting from the proposed approach is better and has a better extrapolation ability.

5 Conclusion

In various applications (*e.g.* safety/security, finance, insurance, environmental sciences), it is of prime importance to predict the response Y of a system when it is impacted by shocks, corresponding to extremely large input values X . In this paper, we have developed a rigorous probabilistic framework for binary classification in extreme regions, relying on the (nonparametric) theory of regularly varying random vectors, and proved the accuracy of the ERM approach in this context, when the risk functional is computed from extreme observations only. The present contribution may open a new line of research, insofar as progress can be naturally expected in the design of algorithmic learning methods tailored to extreme points (or their projection onto the unit sphere) and statistical issues such as estimation of the minimum risk in the extremes, L_∞^* , remain to be addressed.

References

- [1] C. Brownlees, E. Joly, and G. Lugosi. Empirical risk minimization for heavy-tailed losses. *Ann. Statist.*, 43(6):2507–2536, 2015.
- [2] J.J. Cai, J.H.J. Einmahl, and L. De Haan. Estimation of extreme risk regions under multivariate regular variation. *The Annals of Statistics*, pages 1803–1826, 2011.
- [3] A. Carpentier and M. Valko. Extreme bandits. In *Advances in Neural Information Processing Systems 27*, pages 1089–1097. Curran Associates, Inc., 2014.
- [4] S. Coles and J.A Tawn. Modeling extreme multivariate events. *JR Statist. Soc. B*, 53:377–392, 1991.
- [5] L. De Haan and S. Resnick. On regular variation of probability densities. *Stochastic processes and their applications*, 25:83–93, 1987.
- [6] L. Devroye, L. Györfi, and G. Lugosi. *A Probabilistic Theory of Pattern Recognition*. Applications of mathematics : stochastic modelling and applied probability. U.S. Government Printing Office, 1996.
- [7] N. Goix, A. Sabourin, and S. Cléménçon. Learning the dependence structure of rare events: a non-asymptotic study. In *Conference on Learning Theory*, pages 843–860, 2015.
- [8] N. Goix, A. Sabourin, and S. Cléménçon. Sparse representation of multivariate extremes with applications to anomaly ranking. In *Artificial Intelligence and Statistics*, pages 75–83, 2016.
- [9] N. Goix, A. Sabourin, and S. Cléménçon. Sparse representation of multivariate extremes with applications to anomaly detection. *Journal of Multivariate Analysis*, 161:12–31, 2017.
- [10] S. Mendelson. Learning without concentration for general loss functions. *Probability Theory and Related Fields*, 171(1):459–502, 2018.
- [11] K. Nakai and M. Kanehisa. A knowledge base for predicting protein localization sites in eukaryotic cells. *Genomics*, 14(4):897–911, 1992.
- [12] Mesrob I Ohannessian and Munther A Dahleh. Rare probability estimation under regularly varying heavy tails. In *Conference on Learning Theory*, pages 21–1, 2012.
- [13] S. Resnick. *Extreme Values, Regular Variation, and Point Processes*. Springer Series in Operations Research and Financial Engineering, 1987.
- [14] Teemu Roos, Peter Grünwald, Petri Myllymäki, and Henry Tirri. Generalization to unseen cases. In Y. Weiss, B. Schölkopf, and J. C. Platt, editors, *Advances in Neural Information Processing Systems 18*, pages 1129–1136. MIT Press, 2006.
- [15] A. Stephenson. Simulating multivariate extreme value distributions of logistic type. *Extremes*, 6(1):49–59, 2003.

APPENDIX

A Regular Variation and vague convergence

A Borelian function $U : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is said to be regularly varying at infinity if there exists a function $h(x)$ such that

$$\lim_{t \rightarrow \infty} \frac{U(tx)}{U(t)} = h(x), \quad \forall x > 0.$$

In such a case the function h is necessarily of the form $h(x) = x^\rho$ for some $\rho \in \mathbb{R}$. We denote $U \in \mathcal{RV}_\rho$. Now let X be a random variable with *cdf* F and survival function $\bar{F} = 1 - F$. The random variable X is said to be heavy tailed with tail index $\alpha > 0$ if $\bar{F} \in \mathcal{RV}_{-\alpha}$. In addition, the heavy-tail property can be classically formulated in terms of vague convergence to a homogeneous Radon measure. Indeed, the random variable X is heavy-tailed with index α if and only if there exists a sequence $b_n > 0$ with $b_n \rightarrow \infty$, such that

$$n\mathbb{P}\{X/b_n \in \cdot\} \xrightarrow[n \rightarrow \infty]{v} \mu_\alpha(\cdot) \text{ in } M_+(0, \infty],$$

where \xrightarrow{v} denotes vague convergence, $\mu_\alpha(x, \infty]/\mu_\alpha(1, \infty] = x^{-\alpha}$ and $M_+(0, \infty]$ is the set of nonnegative Radon measures on $[0, \infty] \setminus \{0\}$. Then the sequence b_n may be chosen as $b_n = F^{-1}(1 - 1/n)$, where $F^{-1}(u) = \inf\{t : F(t) \geq u\}$ denotes F 's generalized inverse, or $b_n = n^{1/\alpha}$.

We recall that vague convergence of a sequence of Radon measures ν_n on a Locally Compact, Second Countable, Hausdorff space E is defined in terms of convergence of integrals of continuous functions with compact support. Namely, $\nu_n \xrightarrow{v} \nu$ in $M_+(E)$ if and only if for any such function f , $\int_E f d\nu_n \xrightarrow[n \rightarrow \infty]{} \int_E f d\nu$. The statements of Section 2.1 are based on following version of the Portmanteau theorem for vague convergence: $\nu_n \xrightarrow{v} \nu$ if and only if for all relatively compact set $A \subset E$ such that $\nu(\partial A) = 0$, $\nu_n(A) \xrightarrow[n \rightarrow \infty]{} \nu(A)$.

B Technical proofs

Proof of Theorem 1

Letting P denote the law of X , in view of expression (6) for $L_{P_\infty}^*$,

$$\begin{aligned} L_t(g^*) - L_{P_\infty}^* &= L_t(g^*) - \mathbb{E}[\min\{\eta_\infty(X_\infty), 1 - \eta_\infty(X_\infty)\}] \\ &\leq \frac{t \int_{\{\|x\| > t\}} \min\{\eta(x), 1 - \eta(x)\} - \min\{\eta_\infty(x), 1 - \eta_\infty(x)\} dP(x)}{t\mathbb{P}\{\|X\| > t\}} \\ &\quad + \left(\frac{t \int_{\{\|x\| > t\}} \min(\eta_\infty, 1 - \eta_\infty) dP}{t\mathbb{P}\{\|X\| > t\}} - \frac{\int_{\{\|x\| > 1\}} \min(\eta_\infty, 1 - \eta_\infty) d\mu}{\mu(\Omega^c)} \right) \\ &:= A + B \end{aligned}$$

The first term is controlled by Assumption 2. Indeed

$$A \leq \sup_{r \geq t} \sup_{\theta \in S} |\eta(r\theta) - \eta_\infty(\theta)|$$

which goes to 0 as $t \rightarrow \infty$ under Assumption 2. Now regular variation of F means that for any continuous function h with compact support in $[0, \infty]^d \setminus \{0\}$, (that is with support bounded away from 0), $t\mathbb{E}\{h(t^{-1}X)\} \rightarrow \int h d\mu$, which implies, using the continuity assumption on η_∞ , that $B \rightarrow 0$ as well.

We now turn to the second assertion of the theorem. Since $L_\infty^* = L_\infty(g^*)$, any classifier \tilde{g} such that

$$\limsup_{t \rightarrow \infty} \{L_t(\tilde{g}) - L_t(g^*)\} = 0 \tag{11}$$

minimizes L_∞ as well. We shall thus prove that (11) holds for $\tilde{g} = g_\infty^*$. For any classifier of the kind $g(x) = 2\mathbb{1}\{s(x) > 1/2\} - 1$ where s is a scoring function, we have

$$\begin{aligned} L_t(g) &= \frac{\int_{\{\|x\| > t\}} \eta(x)\mathbb{1}\{s(x) < 1/2\} + (1 - \eta(x))\mathbb{1}\{s(x) > 1/2\} dP(x)}{\mathbb{P}\{\|X\| > t\}} \\ &= \mathbb{E}\{(2\eta(X) - 1)\mathbb{1}\{s(X) < 1/2\} \mid \|X\| > t\} + \mathbb{E}\{1 - \eta(X) \mid \|X\| > t\}, \end{aligned}$$

thus

$$L_t(g_\infty^*) - L_t(g^*) = \mathbb{E} \{ (2\eta(X) - 1) (\mathbf{1}\{\eta_\infty(X) < 1/2\} - \mathbf{1}\{\eta(X) < 1/2\}) \mid \|X\| > t \}.$$

Let $0 < \epsilon < 1/2$. We may write

$$L_t(g_\infty^*) - L_t(g^*) = \frac{1}{t\mathbb{P}\{\|x\| > t\}} (A + B + C),$$

with

$$\begin{aligned} A &= t \int_{\|x\| > t, \eta_\infty(x) < 1/2 - \epsilon} (2\eta(x) - 1) (1 - \mathbf{1}\{\eta(x) < 1/2\}) \, dP(x), \\ B &= t \int_{\|x\| > t, \eta_\infty(x) > 1/2 + \epsilon} (2\eta(x) - 1) (-\mathbf{1}\{\eta(x) < 1/2\}) \, dP(x), \\ C &= t \int_{\|x\| > t, |\eta_\infty(x) - 1/2| \leq \epsilon} (2\eta(x) - 1) (\mathbf{1}\{\eta_\infty(x) < 1/2\} - \mathbf{1}\{\eta(x) < 1/2\}) \, dP(x). \end{aligned}$$

For $t_0 > 0$ such that $\sup_{\|x\| > t_0} |\eta(x) - \eta_\infty(x)| < \epsilon/2$ (see Remark 4), the integrands in A and B are zero. On the other hand,

$$|C| \leq 2\epsilon * 2t\mathbb{P}\{\|X\| > t\}$$

Thus for $t > t_0$, $L_t(g_\infty^*) - L_t(g^*) < 4\epsilon$. Since ϵ is arbitrarily small, the proof is complete.

Proof of Theorem 2

The proof relies on the classical bias/variance risk decomposition which takes the following form for the risk above level t :

$$L_{t_\tau} - L_{t_\tau}^* \leq 2 \sup_{g \in \mathcal{G}_S} |\widehat{L}_k(g) - L_{t_\tau}(g)| + \inf_{g \in \mathcal{G}_S} L_{t_\tau}(g) - L_{t_\tau}^*. \quad (12)$$

The statement of the theorem then immediately derives from the uniform bound on the deviations of \widehat{L}_k of the class \mathcal{G}_S stated in Theorem 3 below.

Theorem 3 *In the setting of Theorem 2, for all $\delta \in (0, 1)$, we have with probability $1 - \delta$:*

$$\begin{aligned} \sup_{g \in \mathcal{G}_S} |\widehat{L}_k(g) - L_{t_\tau}(g)| &\leq \frac{1}{\sqrt{k}} \left(\sqrt{2(1 - \tau) \log(2/\delta)} + C \sqrt{V_{\mathcal{G}_S} \log(1/\delta)} \right) \\ &\quad + \frac{1}{k} \left(5 + 2 \log(1/\delta) + \sqrt{\log(1/\delta)} (C \sqrt{V_{\mathcal{G}_S}} + \sqrt{2}) \right) \end{aligned}$$

where C is a constant independent of n, τ, δ .

PROOF. [Proof of Theorem 3] Set $k = \lfloor n\tau \rfloor$ throughout. Introduce the pseudo-empirical risk

$$\widetilde{L}_k = \frac{1}{k} \sum_{i=1}^n \mathbf{1}\{g(X_i) \neq Y_i, \|X_i\| \geq t_\tau\}.$$

Notice that \widetilde{L}_k is not observed since t_τ is unknown: it serves a useful intermediate quantity in the following excess risk decomposition:

$$\sup_{g \in \mathcal{G}_S} |\widehat{L}_k(g) - L_{t_\tau}(g)| \leq \underbrace{\sup_{g \in \mathcal{G}_S} |\widehat{L}_k(g) - \widetilde{L}_k(g)|}_A + \underbrace{\sup_{g \in \mathcal{G}_S} |\widetilde{L}_k(g) - L_{t_\tau}(g)|}_B \quad (13)$$

The remainder of the proof consists in controlling the first term A in the r.h.s. of (13) via the Bernstein inequality while the second term B requires a call to a VC inequality for low probability

regions. As for the first term,

$$\begin{aligned}
A &\leq \frac{1}{k} \left| \sum_1^n \mathbf{1}\{g(X_i) \neq Y_i\} (\mathbf{1}\{\|X_i\| \geq X_{(k)}\} - \mathbf{1}\{\|X_i\| \geq t_\tau\}) \right| \\
&\leq \frac{1}{k} \sum_1^n |\mathbf{1}\{\|X_i\| \geq \|X_{(k)}\|\} - \mathbf{1}\{\|X_i\| \geq t_\tau\}| \\
&= \frac{1}{k} \sum_1^k |1 - \mathbf{1}\{\|X_{(i)}\| \geq t_\tau\}| + \frac{1}{k} \sum_{k+1}^n |\mathbf{1}\{\|X_{(i)}\| \geq t_\tau\}| \\
&= \begin{cases} \frac{1}{k} \sum_{k+1}^n \mathbf{1}\{\|X_{(i)}\| \geq t_\tau\} & \text{if } \|X_{(k)}\| \geq t_\tau \\ \frac{1}{k} \sum_1^k \mathbf{1}\{\|X_{(i)}\| < t_\tau\} & \text{otherwise} \end{cases} \\
&= \begin{cases} \frac{1}{k} \sum_1^n \mathbf{1}\{\|X_{(i)}\| \geq t_\tau\} - \frac{k}{k} & \text{if } \|X_{(k)}\| \geq t_\tau \\ \frac{1}{k} \sum_1^n \mathbf{1}\{\|X_{(i)}\| < t_\tau\} - \frac{n-k}{k} & \text{otherwise} \end{cases} \\
&= \left| \frac{1}{k} \sum_1^n \mathbf{1}\{\|X_i\| \geq t_\tau\} - 1 \right| \\
&\leq \frac{|S_n - n\tau|}{k} + \frac{1}{k}
\end{aligned}$$

where $S_n = \sum_1^n W_i$ and $W_i = \mathbf{1}\{\|X_i\| > t_\tau\}$. Since $\mathbb{E}\{W_i\} = n\tau$, Bernstein inequality implies, for $y > 0$, $\mathbb{P}\{|S_n - n\tau|/k > y\} \leq 2 \exp\{-y^2 k^2 / 2 / (n\tau(1-\tau) + yk/3)\} := \delta$. Solving the latter bound for y yields $y = \frac{1}{k} \left(\frac{1}{3} \log(2/\delta) + \sqrt{(1/3 \log(2/\delta))^2 + 2n\tau(1-\tau) \log(2/\delta)} \right)$. Simplifying the latter bound using that for $a, b > 0$, $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$, and that $n\tau \leq k+1$, we obtain that with probability $1 - \delta$,

$$A \leq \sqrt{\frac{2}{k}(1-\tau) \log(2/\delta)} + \frac{1}{k} \left(\frac{2}{3} \log(2/\delta) + \sqrt{2(1-\tau) \log(2/\delta)} + 1 \right) \quad (14)$$

We now turn to the second term (B) in (13). Write

$$\begin{aligned}
B &\leq \frac{n}{k} \sup_{g \in \mathcal{G}_S} \underbrace{\left| \frac{1}{n} \sum \mathbf{1}\{g(X_i) \neq Y_i, \|X_i\| \geq t_\tau\} - \mathbb{P}\{g(X) \neq Y, \|X\| \geq t_\tau\} \right|}_{B_1} \\
&\quad + \underbrace{\left| \frac{1}{\tau} - \frac{n}{k} \right| \mathbb{P}\{g(X) \neq Y, \|X\| \geq t_\tau\}}_{B_2}
\end{aligned}$$

First,

$$B_2 \leq \left| \frac{1}{\tau} - \frac{n}{k} \right| \tau = |(k - n\tau)/k| \leq 1/k. \quad (15)$$

Turning to B_1 , from Theorem 1 in [7], we have that for any class of sets \mathcal{A} on an input space \mathcal{Z} with finite VC dimension $V_{\mathcal{A}}$, if $(Z_i)_{i \leq n}$ are *i.i.d.* copies of a r.v. Z , then

$$\sup_{A \in \mathcal{A}} \left| \mathbb{P}\{Z \in A\} - \frac{1}{n} \sum_{i=1}^n \mathbf{1}\{Z_i \in A\} \right| \leq C \left(\sqrt{p} \sqrt{\frac{V_{\mathcal{A}}}{n} \log \frac{1}{\delta}} + \frac{1}{n} \log \frac{1}{\delta} \right) \quad (16)$$

where $p = \mathbb{P}\{Z \in \bigcup_{A \in \mathcal{A}} A\}$ and C is an absolute constant.

By setting $Z = (X, Y)$, $\mathcal{A} = \{\mathcal{A}_g : g \in \mathcal{G}_S\}$ where $\mathcal{A}_g = \{(x, y), g(x) \neq y, \|x\| > t_\tau\}$, so that

$p = \tau$, Equation (16) becomes:

$$\begin{aligned}
B_2 &\leq \frac{n}{k} \sup_{g \in \mathcal{G}_S} \left| \mathbb{P}\{Z \in A_g\} - \frac{1}{n} \sum \mathbb{1}\{Z_i \in A_g\} \right| \\
&\leq \frac{n}{k} C \left(\sqrt{\tau} \sqrt{\frac{V_{\mathcal{G}_S}}{n} \log(1/\delta)} + 1/n \log(1/\delta) \right) \\
&= C \left(\sqrt{\frac{V_{\mathcal{G}_S} n \tau}{k^2} \log(1/\delta)} + \frac{1}{k} \log(1/\delta) \right) \\
&\leq C \left(\sqrt{\frac{V_{\mathcal{G}_S}}{k} \log(1/\delta)} + \frac{1}{k} \left(\log(1/\delta) + \sqrt{V_{\mathcal{G}_S} \log(1/\delta)} \right) \right) \tag{17}
\end{aligned}$$

Combining equations (14), (17) and (15), we obtain

$$\begin{aligned}
\sup_{g \in \mathcal{G}_S} |\widehat{L}_k(g) - L_{t_\tau}(g)| &\leq \sqrt{\frac{2}{k}(1-\tau) \log(2/\delta)} + \frac{1}{k} \left(\frac{2}{3} \log(2/\delta) + \sqrt{2(1-\tau) \log(2/\delta)} + 1 \right) \\
&\quad + \frac{1}{k} + C \left(\sqrt{\frac{V_{\mathcal{G}_S}}{k} \log(1/\delta)} + \frac{1}{k} \left(\log(1/\delta) + \sqrt{V_{\mathcal{G}_S} \log(1/\delta)} \right) \right) \\
&\leq \frac{1}{\sqrt{k}} \left(\sqrt{2(1-\tau) \log(2/\delta)} C \sqrt{V_{\mathcal{G}_S} \log(1/\delta)} \right) \\
&\quad + \frac{1}{k} \left(5 + 2 \log(1/\delta) + \sqrt{\log(1/\delta)} (C \sqrt{V_{\mathcal{G}_S}} + \sqrt{2}) \right)
\end{aligned}$$

■

C Numerical experiments

C.1 Synthetic data from the Clover distribution

The Clover distribution introduced in [2] has the following density

$$f(x, y) = \begin{cases} \frac{3}{10\pi} r_0^4 (1 + r_0^6)^{-\frac{3}{2}} \left(5 + \frac{4(x^2 + y^2)^2 - 32x^2 y^2}{r_0(x^2 + y^2)^{-\frac{3}{2}}} \right) & x^2 + y^2 < r_0 \\ \frac{3}{10\pi} \left(\frac{9(x^2 + y^2)^2 - 32x^2 y^2}{r_0(x^2 + y^2)^{-\frac{3}{2}}} \right) & x^2 + y^2 \geq r_0 \end{cases} \quad \text{with } r_0 = 1.248$$

Without loss of generality, only the points within the positive orthant are kept. Points labeled -1 are rotated by an angle θ in the counterclockwise direction. Figure 6 provides an example of 2D points from both distributions with $\theta = \frac{\pi}{4}$. the number of trees for both random forests (in the regular setting and in the setting of Algorithm 1) is set to 200. The number of neighbors for both k-NN's is set to 5.




Figure 6: Labeled dataset generated from a Clover distribution and its θ -rotated version.

Figures 7 and 8 illustrate once again the conclusions from Section 4.




Figure 7: Clover data - test loss of random forest on the simplex and regular random forest depending on the multiplicative factor κ .




Figure 8: Clover data - test loss of k-NN on the simplex and regular k-NN depending on the multiplicative factor κ .

C.2 Synthetic data from the Logistic distribution

The multivariate logistic model is a widely used model in the context of extreme value analysis, see *e.g.* [4] or [15] for efficient simulation algorithms. The logistic distribution in \mathbb{R}^d with parameter $\delta \in (0, 1]$ has cumulative distribution function

$$F(x) = \exp \left\{ - \left(\sum_{i=1}^d x_j^{\frac{-1}{\delta}} \right)^\delta \right\}, \quad x \in (0, \infty)^d.$$

For small values of δ , extremes tend to occur simultaneously in all direction, that is, the angular measure concentrates around the center of the positive orthant of the unit sphere. On the other hand, for δ close to 1, extremes tend to concentrate near the axes, that is only one component at a time is likely to be large. The positive and negative instances are respectively generated according to a logistic distribution with parameter $\delta_+ = 0.1$ and $\delta_- = 0.5$, in dimension $d = 4$. Ten datasets are simulated. Each one is composed of a train set and a test set containing respectively 10^4 and 10^5 instances. k the number of points used for training the classifier in Algorithm 1) is set to 100. The multiplicative factor κ is made to vary between 1 and 0.3. Figure 9 displays evolutions of boxplots of Hamming losses depending on the multiplicative factor κ : each box summarizes the distribution of the losses obtained with the 10 considered datasets. It is analogue to Figure 4 using k-NN instead of RF, from which similar conclusions can be drawn.




Figure 9: Logistic data - test loss of k-NN on the simplex and regular k-NN depending on the multiplicative factor κ .

C.3 Real-world data: Ecoli dataset

The Ecoli dataset contains 336 instances. There are 8 features: 7 numerical features and 1 corresponding to a sequence name. In our experimental framework the feature corresponding to the

sequence name is dropped: we only work with the other features. The label to be predicted corresponds to the protein localization site among the 8 different localizations possible. In our experiments the labeling is simplified: data labeled *im* are set to 1, all instances labeled differently are set to -1 . The classification problem is thus turned into a binary one. Figure 10 is the analogue of Figure 5 using k-NN instead of RF, from which similar conclusions can be drawn.




Figure 10: Real data - test loss of regular k-NN depending on the multiplicative factor κ .

Remark 7 (On the truncation step) In all experiments, the value of k is set to 100. Finding k , the number of extreme observations involved in the computation of the empirical risk (9), is beyond the scope of this paper though one could expect better results by improving the selection of k .